

## ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
0.67ml SEPP® APPLICATOR

**WARNING, FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

### Drug Facts

**Active ingredients**  
Chlorhexidine gluconate 2% w/v..... Antiseptic  
Isopropyl alcohol 70% v/v..... Antiseptic

**Use** for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.

#### Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

#### Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges.
- on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm<sup>2</sup>)
- remove applicator from package; do not touch applicator tip
- hold the sponge tip down and pinch the barrel of the applicator **only once** to activate the ampule and release the antiseptic
- wet the sponge tip by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information** ■ store between 15–30 °C (59–86 °F)  
■ avoid freezing and excessive heat above 40 °C (104 °F)

**Inactive ingredients** ■ USP purified water

**Questions?** ■ [www.chloroprep.com](http://www.chloroprep.com)  
■ call 1-800-523-6502 (M-F 9 a.m.–5 p.m. CST)

See other side for additional information

## ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
0.67ml SEPP® APPLICATOR

**WARNING, FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

### ⚠ WARNING FLAMMABLE



Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

See Drug Facts for Full Labeling Information



Do Not Reuse



Labex Free

External Use Only

Professional Use Only

Single Use

CareFusion Leawood, KS 66211

Artwork 6-775449F • L10312

Made in the USA of US and imported parts