

# The finishing touch for cutting-edge surgeries

Mepilex® Border Post-Op Ag for minimally invasive surgical incisions



9x10 cm actual size



6x8 cm actual size

You've invested so much in minimally invasive procedures.  
**Choose a post-op dressing worthy of your work.**

Mepilex Border Post-Op Ag is a cut above ordinary tape and gauze. In fact, its strong clinical evidence has helped make the dressing a popular choice across all surgical specialties.

Now, with new **6x8 cm and 9x10 cm sizes added to the Mepilex Border Post-Op Ag portfolio**, you can apply the benefits of our best-in-class dressing to your smallest surgical incisions.

Mepilex® Border Post-Op | Mepilex® Border Post-Op Ag

  
Mölnlycke®



## Gentle on the peri-wound skin

Made with our Safetac® wound contact layer, the dressing adheres firmly but gently to the patient's skin. This minimizes skin damage and pain during dressing removal.<sup>1,2</sup> Anticipatory pain affects some patients<sup>3</sup> and Safetac can help lessen their anxiety.



## Allows both showering and movement

With its showerproof seal and bacteria barrier (microbes larger than 25 nm),<sup>4</sup> patients can feel confident getting Mepilex® Border Post-Op Ag wet. Meanwhile, innovative Flex technology keeps the dressing soft, comfortable and conformable to movement. Perfect for those hips and knees.<sup>2,5,6,7,8</sup>



## Reinforces low SSI rates

While minimally invasive surgeries already pose lower infection risk, Mepilex Border Post-Op Ag adds a layer of protection by starting to inactivate pathogenic bacteria within 30 minutes of application.<sup>9</sup> It continues to inactivate 99.99% of common microbial threats throughout the 7-day<sup>9</sup> dressing wear time.



## No post-surgery cutting

Clinicians can save time in the operating room (and potentially reduce product waste) by not having to work with tape and gauze. Mepilex Border Post-Op Ag is easy to apply. There's nothing to cut.

Minimally invasive procedures elevate the patient experience.  
**Now there's a post-op dressing to match.**

### Mepilex® Border Post-Op

Product code	Dressing size inches (cm)	Pad size inches (cm)	Pcs Bx/Cs
496100	2.5"x3" (6x8 cm)	1"x2" (3x5 cm)	10/100
496200	3.5"x4" (9x10 cm)	2"x2.5" (5x6 cm)	10/100
496300	4"x6" (10x15 cm)	2"x4" (5x10 cm)	10/100
496405	4"x8" (10x20 cm)	2"x6" (5x15 cm)	5/25
496455	4"x10" (10x25 cm)	2"x8" (5x20 cm)	5/30
496605	4"x12" (10x30 cm)	2"x10" (5x25 cm)	5/25
496650	4"x14" (10x35 cm)	2"x12" (5x30 cm)	5/55

### Mepilex® Border Post-Op Ag

Product code	Dressing size inches (cm)	Pad size inches (cm)	Pcs Bx/Cs
498100	2.5"x3" (6x8 cm)	1"x2" (3x5 cm)	5/20
498200	3.5"x4" (9x10 cm)	2"x2.5" (5x6 cm)	5/50
498300	4"x6" (10x15 cm)	2"x4" (5x10 cm)	5/70
498400	4"x8" (10x20 cm)	2"x6" (5x15 cm)	5/25
498450	4"x10" (10x25 cm)	2"x8" (5x20 cm)	5/35
498600	4"x12" (10x30 cm)	2"x10" (5x25 cm)	5/25
498650	4"x14" (10x35 cm)	2"x12" (5x30 cm)	5/60

## Try Mepilex Border Post-Op Ag on your patients.

We cover the full range of incision sizes:  
1 centimeter to 14 inches.

## For a trial, contact your Mölnlycke representative.

[molnlycke.us/products-solutions/mepilex-border-post-op](https://molnlycke.us/products-solutions/mepilex-border-post-op)  
+1 (800) 882-4582

1. V Overschelde P. et al. A randomised controlled trial comparing two wound dressings used after elective hip and knee arthroplasty. Poster presentation at the 5th WUWHS conference, Florence, Italy, 2016. 2. Bredow J. et al. Evaluation of Absorbent Versus Conventional Wound Dressing. A Randomized Controlled Study in Orthopedic Surgery. Deutsche Arzteblatt International, 2018. 3. Upton, Dominic. "Psychological aspects of wound care: implications for clinical practice." Journal of Community Nursing, vol. 28, no. 2, Apr.-May 2014, pp. 52+. Gale Academic OneFile, link.gale.com/apps/doc/A393988700/AONE?u=anon-e6a01e91&sid=googleScholar&xid=22336ad4; Accessed 21 Dec. 2022. 4. Mölnlycke Health Care, Data on file: Viral penetration test. 5. Zarghooni, K. et al. Is the use of modern versus conventional wound dressings warranted after primary knee and hip arthroplasty? Acta Orthopaedica Belgica, 2015. 6. Van Overschelde P. et al. A randomised controlled trial comparing two wound dressings used after elective hip and knee arthroplasty. Poster presentation at 5th Congress of WUWHS, Florence, Italy 2016. 7. Johansson C et al. Preventing post-operative blisters following hip and knee arthroplasty. Wounds International, 2012. 8. Korby K. et al. Evidence-Based Practice: Clinical evaluation of a new silver soft silicone postoperative dressing in total knee and hip arthroplasty. Poster presentation at the PAOS conference, US, 2017. 9. Mölnlycke Health Care, Data on file. Laboratory tests 20151026-005, 20151109-002, 20151110-007. The Mölnlycke® and Mepilex® trademarks, names and logo types are registered globally to one or more of the Mölnlycke Health Care Group of Companies. Distributed by Mölnlycke Health Care US, LLC, Peachtree Corners, Georgia 30092. © 2023 Mölnlycke Health Care AB. All rights reserved. 1-800-843-849 MHC-2023-110130

